

## Travel Nurse HAI Prevention Guide

Thank you for joining our team!

Here are a few tips to help us prevent Hospital Acquired Infections (HAIs) in our patients.

### CLABSI

- ☐ Perform **Hand Hygiene** before and after touching a Central line. Assist patient in performing hand hygiene frequently.
- ☐ **Change dressing** every 7 days unless wet, loose, or soiled, then change as soon as possible. Dressing changes are done on Thursdays unless needed.
  - If using gauze under dressing, change every 48 hrs.
- ☐ Keep **alcohol caps** on ports when not in use. Use an alcohol scrub when alcohol caps aren't being used, if port becomes contaminated, and with each sequential access.
  - Use a new cap each time one is removed.
- ☐ **Change needleless connectors** every 96 hours or when occluded or soiled.
- ☐ **Flush** line every 12 hours when not in use.
- ☐ **Document** blood return every 12 hours.
- ☐ Daily **bathing** is important. **CHG treatment after bathing** required for all patients in ICUs and for patients with Central Lines on acute care.
- ☐ When drawing a blood culture, **peripheral sticks are preferred**. Do not draw blood cultures from Central Lines.

### CAUTI

- ☐ Perform **Hand Hygiene** before and after touching a Foley Catheter. Assist patient in performing hand hygiene frequently.
- ☐ Keep foley **below the bladder** during transports and when ambulating.
- ☐ **Perform peri- cares with Provon wipes** every shift and after incontinence or stooling.
- ☐ Keep foley **secure with a Stat-lock** to prevent pistoning.
- ☐ **Maintain a closed system** (keep tamper seal intact).
- ☐ Use **two caregivers during insertion** to maintain sterile field.
- ☐ **Remove foley** as soon as possible when no longer clinically indicated.
- ☐ **Perform daily bathing** to reduce bioburden.
- ☐ **Check with the Charge nurse** before sending a urine sample to assure the correct steps are taken for collection.

## CDI

- ☐ Perform **Hand Hygiene** before and after any cares. Assist patient in performing hand hygiene frequently. After a Cdiff diagnosis, Hand hygiene should be performed with soap and water.
- ☐ **Perform daily bathing** to reduce bioburden.
- ☐ **High touch cleaning** should be performed every shift. After a C-diff diagnosis, high touch cleaning should be performed with bleach.
- ☐ If the patient has **diarrhea on admit**, ask the doctor about sending a C-diff sample **before hospital day 4**.
- ☐ **Check with the Charge nurse** before sending a C-diff sample to ensure the correct steps are taken for collection and that the test has been ordered appropriately.



**Please document all interventions!**

### About this Job Aid |

**Author(s):** Bonnie St. John, Infection Prevention and Control Professional II-Non-Exempt

**Approved by:** Melodie Toll, Associate CNO, Specialty Based Care

**Learning Network Contact:** Joyce A Gamble Clinical Education Director

**Expiration Date:** 11/30/2022